

ERAs for menopause treatment: Welcome the 'designer estrogens'

Cleveland Clinic Journal of Medicine. 2017 June;84(6):463-470



TAMOXIFEN: CANCER TREATMENT AND PREVENTION

After clomiphene, tamoxifen was the second ERAA on the market. Although researchers were looking for a new contraceptive drug, they found tamoxifen to be useful as a chemotherapeutic agent for breast cancer. First used in 1971, tamoxifen continues to be one of the most commonly prescribed chemotherapeutic medications today.

The FDA has approved tamoxifen to treat breast cancer as well as to prevent breast cancer in pre- and postmenopausal women at risk. It may also have beneficial effects on bone and on cardiovascular risk factors, but these are not approved uses for it.

Trials of tamoxifen for cancer treatment

The Early Breast Cancer Trialists' Collaborative Group⁹ performed a meta-analysis and found that 5 years of adjuvant treatment with tamoxifen is associated with a 26% reduction in mortality and a 47% reduction in breast cancer recurrence at 10 years. In absolute terms, we estimate that 21 women would need to be treated to prevent 1 death and 8 would need to be treated to prevent 1 recurrence.

The ATLAS Trial (Adjuvant Tamoxifen Longer Against Shorter)¹⁰ and later the UK **ATTOM** (Adjuvant Tamoxifen Treatment to Offer More)¹¹ trial confirmed an even greater reduction in recurrence and mortality after a total of 10 years of treatment.

Trials of tamoxifen for cancer prevention

Cuzik et al¹² performed a meta-analysis of 4 trials of tamoxifen's effectiveness in

preventing breast cancer for women at elevated risk. The incidence of estrogen receptor-positive breast cancer was 48% lower with tamoxifen use, but there was no effect on estrogen-negative breast cancer. From their data, we estimate that 77 women would need to be treated to prevent 1 case of breast cancer.

The IBIS-I trial (International Breast Cancer Intervention Study I)¹³ found that, in healthy women at high risk of breast cancer, the benefit of taking tamoxifen for 5 years as preventive treatment persisted long afterward. The investigators estimated that at 20 years of follow-up the risk of breast cancer would be 12.3% in placebo recipients and 7.8% in tamoxifen recipients, a 4.5% absolute risk reduction; number needed to treat (NNT) 22.

Data on tamoxifen and osteoporosis

The Breast Cancer Prevention Trial revealed a 19% reduction in the incidence of osteoporotic fractures with tamoxifen, but the difference was not statistically significant.¹⁴ The 1-year rates of fracture in women age 50 and older were 0.727% with placebo and 0.567% with tamoxifen, an absolute difference of 0.151%; therefore, if the effect is real, 662 women age 50 or older would need to be treated for 1 year to prevent 1 fracture. Tamoxifen is not FDA-approved to treat osteoporosis.

Data on tamoxifen and cardiovascular risk reduction

Chang et al,¹⁵ in a study in women at risk of breast cancer, incidentally found that tamoxifen was associated with a 13% reduction in total cholesterol compared with placebo.

Herrington and Klein,¹⁶ in a systematic review, noted similar findings in multiple studies of tamoxifen, with decreases in total cholesterol ranging from 7% to 17% and decreases in low-density lipoprotein cholesterol ranging from 10% to 28%. However, they found no change in high-density lipoprotein cholesterol concentrations or in the cardiovascular mortality rate.

The ATLAS trial¹⁰ revealed a relative risk reduction of 0.76 (95% confidence interval [CI] 0.60–0.95, $P = .02$) in ischemic heart disease for women who took tamoxifen for 10 years compared with 5 years. We calculate that ischemic heart disease occurred in 163

(2.5%) of 6,440 women who took tamoxifen for 5 years compared with 127 (1.9%) of 6,454 women who took it for 10 years, a 0.6% absolute risk reduction, NNT = 167.

Adverse effects of tamoxifen

Uterine neoplasia. Women taking tamoxifen have a 2.5-fold increased risk of endometrial cancer.¹⁴ Tamoxifen also increases the risk of benign uterine disease such as endometrial hyperplasia and polyps. As many as 39% of women taking tamoxifen will have evidence of benign uterine changes on pathology.¹⁷ Other adverse effects:

Venous thromboembolism (the risk of pulmonary embolism is increased approximately threefold¹⁴)

Cataracts (there is a slight increase in cataract diagnosis in tamoxifen users)

Vasomotor symptoms, which limit the use of tamoxifen in many women.

Ideal candidate for tamoxifen

The ideal candidate for tamoxifen is a woman with breast cancer that is estrogen receptor-positive and who has a history of osteopenia or osteoporosis and no risk factors for venous thromboembolism.

« first ‹ previous 1 **2** 3 4 next › last »

DRUG THERAPY

ENDOCRINOLOGY

GERIATRICS

ONCOLOGY

PREVENTIVE CARE

WOMEN'S HEALTH

Related Articles

REVIEWS

Is there a time limit for systemic menopausal hormone therapy?

The duration needs to be an individualized, shared decision that is assessed annually.



THIS PUBLICATION

[About CCJM](#), [Submit A Manuscript](#), [Author Information](#), [Subscription Information](#), [CME Calendar](#),
[Advertise With CCJM](#), [RSS Feeds](#), [Contact Us](#)

TOP SECTIONS

[CME](#), [Reviews](#), [1-Minute Consult](#), [Interpreting Key Trials](#), [IM Board Review](#), [Smart Testing](#)

HIGHLIGHTED FROM THIS SITE AND NETWORK

[About Cleveland Clinic](#), [Cleveland Clinic Education Institute](#),
[Cleveland Clinic Center for Continuing Education](#),
[Cleveland Clinic Resources for Medical Professionals](#),
[Cleveland Clinic Consumer Health Information](#), [Consult QD](#), [Contact ClinicalEdge Manager](#),
[Contact MD-IQ Manager](#), [MedJob Network](#)

[Advertise with MDedge](#)

See more with MDedge! [See our Other Publications](#)

MDedge: Your Source for Clinical Advantage

Copyright © 2017 Cleveland Clinic. All rights reserved. The information provided is for educational purposes only. Use of this website is subject to the [disclaimer](#) and [privacy policy](#).

ERAs for menopause treatment: Welcome the 'designer estrogens'

Cleveland Clinic Journal of Medicine. 2017 June;84(6):463-470



RALOXIFENE: FOR OSTEOPOROSIS AND FOR CANCER PREVENTION

Raloxifene, a second-generation ERAA, was first approved for preventing and treating osteoporosis and later for reducing the risk of invasive estrogen receptor-positive breast cancer in postmenopausal women.

Trials of raloxifene for osteoporosis

The MORE trial (Multiple Outcomes of Raloxifene)¹⁸ was a large multicenter randomized double-blind study. Raloxifene recipients showed a significant increase in bone mineral density in the lumbar spine and femoral neck at year 3 ($P < .001$) compared with those receiving placebo. Even after only 1 year of treatment, raloxifene significantly reduced the risk of new fractures, despite only modest gains in bone mineral density. After 3 years of treatment, new clinical vertebral fractures had occurred in 3.5% of the placebo group compared with 2.1% of the group receiving raloxifene 60 mg.¹⁹ Relative risk reductions were similar in women who had already had a clinical vertebral fracture at baseline, whose absolute risk is higher. However, no significant effect was seen on the incidence of hip or nonvertebral fractures.

The CORE trial (Continuing Outcomes Relevant to Raloxifene)²⁰ extended the treatment of the women enrolled in the MORE trial another 4 years and found that the benefit of raloxifene with regard to bone mineral density persisted with continued use.

Trials of raloxifene for breast cancer prevention

The MORE trial,²¹ in postmenopausal women with osteoporosis included breast cancer as a secondary end point, and raloxifene was shown to decrease the incidence of invasive breast cancer. At a median of 40 months, invasive breast cancer had arisen in 13 (0.25%) of the 5,129 women assigned to raloxifene and 27 (1.0%) of the 2,576 women assigned to placebo. The authors calculated that 126 women would need to be treated to prevent 1 case of breast cancer.

The CORE trial,²² as noted, extended the treatment of the women enrolled in the MORE trial another 4 years. The risk of any invasive breast cancer in postmenopausal women with osteoporosis was significantly reduced by 59% after 8 years, and the risk of estrogen receptor-positive invasive breast cancer was reduced by 66%.

There is evidence that raloxifene's effect on breast cancer risk persists after discontinuation of use.²³

Does raloxifene reduce mortality?

Grady et al²⁴ studied the effect of raloxifene on all-cause mortality in a pooled analysis of mortality data from the MORE, CORE, and Raloxifene Use for the Heart (RUTH)²⁵ trials. In older postmenopausal women, the rate of all-cause mortality was 8.65% in those taking placebo compared with 7.88% in those taking raloxifene 60 mg daily—10% lower. The mechanism behind the lower mortality rate is unclear, and Grady et al recommend that the finding be interpreted with caution.

Trials of raloxifene for heart protection

The RUTH trial²⁵ was a 5.6-year study undertaken to study the effects of raloxifene on coronary outcomes and invasive breast cancer in postmenopausal women. Results were mixed. Active treatment:

- Did not significantly affect the risk of coronary artery disease compared with placebo
- Significantly decreased the risk of invasive breast cancer
- Significantly decreased the risk of clinical vertebral fractures
- Increased the risk of fatal stroke (59 vs 39 events, hazard ratio 1.49, 95% CI 1.00–

2.24) and venous thromboembolism (103 vs 71 events, hazard ratio 1.44, 95% CI 1.06–1.95).

The STAR trial (Study of Tamoxifen and Raloxifene)^{26,27} compared raloxifene and tamoxifen in postmenopausal women at increased risk of breast cancer. Women were randomized to receive either tamoxifen 20 mg or raloxifene 60 mg for 5 years. Results:

- No difference in the number of new cases of invasive breast cancer between the groups
- Fewer cases of noninvasive breast cancer in the tamoxifen group, but the difference was not statistically significant
- Fewer cases of uterine cancer in the raloxifene group, annual incidence rates 0.125% vs 0.199%, absolute risk reduction 0.74%, NNT 1,351, relative risk with raloxifene 0.62, 95% CI 0.30–0.50
- Fewer thromboembolic events with raloxifene
- Fewer cataracts with raloxifene.

Adverse effects of raloxifene

Raloxifene increases the risk of venous thromboembolism and stroke in women at high risk of coronary artery disease.¹⁹

Ideal candidates for raloxifene

Postmenopausal women with osteopenia or osteoporosis and a higher risk of breast cancer who have minimal to no vasomotor symptoms or genitourinary syndrome of menopause are good candidates for raloxifene. Raloxifene is also a good choice for women who have genitourinary syndrome of menopause treated with local vaginal estrogen. Raloxifene has no effect on vasomotor symptoms or genitourinary syndrome of menopause.

Related Articles

REVIEWS

Is there a time limit for systemic menopausal hormone therapy?

The duration needs to be an individualized, shared decision that is assessed annually.



THIS PUBLICATION

[About CCJM](#), [Submit A Manuscript](#), [Author Information](#), [Subscription Information](#), [CME Calendar](#),
[Advertise With CCJM](#), [RSS Feeds](#), [Contact Us](#)

TOP SECTIONS

[CME](#), [Reviews](#), [1-Minute Consult](#), [Interpreting Key Trials](#), [IM Board Review](#), [Smart Testing](#)

HIGHLIGHTED FROM THIS SITE AND NETWORK

[About Cleveland Clinic](#), [Cleveland Clinic Education Institute](#),
[Cleveland Clinic Center for Continuing Education](#),
[Cleveland Clinic Resources for Medical Professionals](#),
[Cleveland Clinic Consumer Health Information](#), [Consult QD](#), [Contact ClinicalEdge Manager](#),
[Contact MD-IQ Manager](#), [MedJob Network](#)

[Advertise with MDedge](#)

See more with MDedge! [See our Other Publications](#)

MDedge: Your Source for Clinical Advantage

Copyright © 2017 Cleveland Clinic. All rights reserved. The information provided is for educational purposes only. Use of this website is subject to the [disclaimer](#) and [privacy policy](#).

ERAAs for menopause treatment: Welcome the 'designer estrogens'

Cleveland Clinic Journal of Medicine. 2017 June;84(6):463-470



OSPEMIFENE: FOR GENITOURINARY SYNDROME OF MENOPAUSE

Although ospemifene does not have the steroid structure of estrogen, it acts as an estrogen agonist specifically in the vaginal mucosa and an antagonist in other tissues.²⁸ It has been shown on Papanicolaou smears to reduce the number of parabasal cells and increase the number of intermediate and superficial cells after 3 months of treatment.²⁹

Ospemifene 60 mg taken orally with food is approved by the FDA to treat genitourinary syndrome of menopause.

Why ospemifene is needed

First-line treatment options for genitourinary syndrome of menopause include over-the-counter lubricants. However, there is no evidence that these products reverse vaginal atrophy,³⁰ and many women report no relief of symptoms with them.

While various local estrogen preparations positively affect genitourinary syndrome of menopause, some of them can be messy, which can limit long-term adherence.

In one of the largest surveys on genitourinary syndrome of menopause (the REVIVE survey—the Real Women's View of Treatment Options for Menopausal Vaginal Changes²⁹), 59% of women reported that their vaginal symptoms negatively affected sexual activity. The problem affects not only the patient but also her sexual partner.³¹ Another large study showed that 38% of women and 39% of male partners reported that it had a worse-than-expected impact on their intimate relationships.³¹

Genitourinary syndrome of menopause also makes pelvic examinations difficult, may worsen or exacerbate cystitis, and may increase urinary tract infections.

Trials of ospemifene for genitourinary syndrome of menopause

To date, 3 randomized, double-blind clinical trials have demonstrated ospemifene 60 mg to be superior to placebo in treating genitourinary syndrome of menopause. Two were short-term (12-week) and showed significant positive changes in the percent of superficial cells, vaginal pH (lower is better), and number of parabasal cells, along with improvements in the Likert rating of both vaginal dryness and dyspareunia.^{32,33}

A long-term (52-week) randomized placebo-controlled trial compared ospemifene and placebo and showed significant improvement in vaginal maturation index and pH at weeks 12 and 52.³⁴ Other outcome measures included petechiae, pallor, friability, erythema, and dryness, all of which improved from baseline ($P < .001$). At the end of the trial, 80% of the patients who received ospemifene had no vaginal atrophy.

No serious adverse events were noted in any of the clinical trials to date, and a systemic review and meta-analysis demonstrated ospemifene to be safe and efficacious.³⁵ The most frequently reported reasons for discontinuation were hot flashes, vaginal discharge, muscle spasms, and hyperhidrosis, but the rates of these effects were similar to those with placebo.

Trial of ospemifene's effect on bone turnover

As an estrogen receptor agonist in bone, ospemifene decreases the levels of bone turnover markers in postmenopausal women.³⁶ A study found ospemifene to be about as effective as raloxifene in suppressing bone turnover,³⁷ but ospemifene does not carry FDA approval for preventing or treating osteoporosis.

Other effects

In experiments in rats, the incidence of breast cancer appears to be lower with ospemifene, and the higher the dose, the lower the incidence.³⁸

Ospemifene also has antagonistic effects on uterine tissue, and no cases of endometrial hyperplasia or carcinoma have been reported in short-term or long-term studies.³⁵

Ospemifene has no effect however on vasomotor symptoms and may in fact worsen vasomotor symptoms in women suffering with hot flashes and night sweats. Further investigation into its long-term safety and effects on breast tissue and bone would provide more insight.

Ideal candidates for ospemifene

Ospemifene could help postmenopausal women with genitourinary syndrome of menopause for whom over-the-counter lubricants fail, who dislike local vaginal estrogen, or who decline systemic hormone therapy, and who do not meet the criteria for treatment with systemic hormone therapy.

CONJUGATED ESTROGENS AND BAZEDOXIFENE COMBINATION

A combination agent consisting of conjugated estrogens 0.45 mg plus bazedoxifene 20 mg has been approved by the FDA for treating moderate to severe vasomotor symptoms associated with menopause and also for preventing postmenopausal osteoporosis in women who have an intact uterus.

Trials of estrogen-bazedoxifene for vasomotor symptoms

The Selective Estrogen Menopause and Response to Therapy (SMART) trials^{39,40} were a series of randomized, double-blind, placebo-controlled phase 3 studies evaluating the efficacy and safety of the estrogen-bazedoxifene combination in postmenopausal women.

The SMART-2 trial³⁹ evaluated the combination of conjugated estrogens (either 0.45 mg or 0.625) plus bazedoxifene 20 mg and found both dosages significantly reduced the number and severity of hot flashes at weeks 4 and 12 ($P < .001$). At week 12, the combination with 0.45 mg of estrogen reduced vasomotor symptoms from baseline by 74% (10.3 hot flashes per week at baseline vs 2.8 at week 12); the combination with 0.625 mg of estrogen reduced vasomotor symptoms by 80% (10.4 vs 2.4 flashes); and placebo reduced them by 51% (10.5 vs 5.4 flashes).

For bone density. The SMART-1 trial⁴⁰ showed that the estrogen-bazedoxifene combination in both estrogen dosages significantly increased mean lumbar spine bone mineral density ($P < .001$) and total hip bone mineral density ($P < .05$) from baseline at

12 and 24 months compared with placebo. Increases in density tended to be higher with the higher estrogen dose (0.625 mg), but less with higher doses of bazedoxifene.⁴¹ At 24 months, the increase in bone mineral density was even greater than in women treated with raloxifene.⁴² However, the effect of estrogen-bazedoxifene on the incidence of fractures remains to be studied.

For genitourinary syndrome of menopause. The SMART-3 trial showed that treatment with conjugated estrogens plus bazedoxifene (0.45/20 mg or 0.625/20 mg) was more effective than placebo in increasing the percent of superficial and intermediate cells and decreased the number of parabasal cells at 12 weeks compared with placebo ($P < .01$).⁴³ Both doses also significantly decreased the mean vaginal pH and improved vaginal dryness.

Patients treated with estrogen-bazedoxifene for a minimum of 12 weeks in a double-blind placebo-controlled study also showed a significant improvement in sexual function and quality-of-life measurements based on 3 well-defined scales, which included ease of lubrication, satisfaction with treatment, control of hot flashes, and sleep parameters.⁴³

Low rates of side effects

To evaluate this regimen's antagonistic effects on uterine tissue, endometrial hyperplasia was diagnosed by blinded pathologists using endometrial biopsies taken at 6, 12, and 24 months or more if cancer was a suspected diagnosis. At 12 and 24 months of treatment, the incidence of hyperplasia with bazedoxifene 20 or 40 mg at doses of either 0.45 or 0.625 mg of conjugated estrogens was less than 1%, which was similar to placebo rates over the 24 months.⁴⁴ The lowest dose studied, bazedoxifene 10 mg, did not prevent hyperplasia with conjugated estrogens 0.45 or 0.625 mg, and its use was discontinued.

Rates of amenorrhea with bazedoxifene 20 or 40 mg and conjugated estrogens 0.45 or 0.625 mg were very favorable (83%–93%) and similar to those with placebo.⁴⁵ For women with continued bleeding on hormone therapy requiring multiple evaluations, or for women who won't accept the risk of bleeding on hormone therapy, conjugated estrogens and bazedoxifene may be a sustainable option. However, any woman with abnormal bleeding should undergo prompt immediate evaluation.

A typical side effect of estrogen replacement therapy is breast tenderness. For women

seeking vasomotor symptom treatment but who experience breast tenderness, this may be a deterrent from continuing hormone therapy. As shown in the SMART-1 and SMART-2 trials,⁴⁶ conjugated estrogens and bazedoxifene did not cause an increase in breast tenderness, which may enhance medication adherence.

Ideal candidates for conjugated estrogens plus bazedoxifene

This product could help postmenopausal women who have an intact uterus and are suffering with moderate to severe vasomotor symptoms and genitourinary syndrome of menopause who cannot tolerate the side effects of hormone therapy such as bleeding, bloating, or breast tenderness, or who prefer to take an estrogen but without a progestin. It is also ideal for women at higher risk of osteoporosis.

WHO SHOULD GET WHAT?

Not all postmenopausal women have vasomotor symptoms, genitourinary syndrome of menopause, or bone loss. For those who do, standard hormone therapy is an option.

For those who have symptoms and a lower threshold of side effects such as breast tenderness and vaginal bleeding, a combination of an estrogen plus an ERAA (eg, bazedoxifene) is an option.

For women who have no vasomotor symptoms but do have genitourinary syndrome of menopause and don't want local vaginal treatment, ospemifene is an option.

For women with no vasomotor symptoms but who have bone loss and increased risk of estrogen receptor-positive breast cancer, raloxifene is a good option.

Both premenopausal and postmenopausal women who are at increased risk for breast cancer should be considered for tamoxifen chemoprevention. Postmenopausal women with a uterus at increased risk for breast cancer should be considered for raloxifene, as it has no uterine effect. Raloxifene is *not* indicated in premenopausal women.

No woman at increased risk of venous thromboembolism is a candidate for ERAA treatment or for oral estrogen. However, the clinician has multiple options to improve quality of life and work productivity and reduce office visits of women at midlife, especially when they are individually assessed and treated.

References

1. Giannini A, Russo E, Mannella P, Simoncini T. Selective steroid receptor modulators in reproductive medicine. *Minerva Ginecol* 2015; 67:431–455.
2. Feldman BM, Voda A, Gronseth E. The prevalence of hot flash and associated variables among perimenopausal women. *Res Nurs Health* 1985; 8:261–268.
3. Versi E, Harvey MA, Cardozo L, Brincat M, Studd JW. Urogenital prolapse and atrophy at menopause: a prevalence study. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; 12:107–110.
4. Hess R, Chang CC, Conigliaro J, McNeil M. Understanding physicians' attitudes towards hormone therapy. *Womens Health Issues* 2005; 15:31–38.
5. Melton LJ 3rd, Khosla S, Atkinson EJ, O'Fallon WM, Riggs BL. Relationship of bone turnover to bone density and fractures. *J Bone Miner Res* 1997; 12:1083–1091.
6. Sikin A, Thacker HL. Treatment options for menopausal hot flashes. *Cleve Clin J Med* 2004; 71:578–582.
7. Levine JP. Treating menopausal symptoms with a tissue-selective estrogen complex. *Gend Med* 2011; 8:57–68.
8. Pinkerton JV, Thomas S. Use of SERMs for treatment in postmenopausal women. *J Steroid Biochem Mol Biol* 2014; 142:142–154.
9. Tamoxifen for early breast cancer: an overview of the randomised trials. Early Breast Cancer Trialists' Collaborative Group. *Lancet* 1998; 351:1451–1467.
10. Davies C, Pan H, Godwin J, et al; Adjuvant Tamoxifen: Longer Against Shorter (ATLAS) Collaborative Group. Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years after diagnosis of oestrogen receptor-positive breast cancer: ATLAS, a randomised trial. *Lancet* 2013; 381:805–816.
11. Gray RG, Rea D, Handley K, et al. aTTom: Long-term effects of continuing adjuvant tamoxifen to 10 years versus stopping at 5 years in 6,953 women with early breast cancer. *J Clin Oncol*

2013; (suppl): abstract 5.

12. Cuzick J, Powles T, Veronesi U, et al. Overview of the main outcomes in breast-cancer prevention trials. *Lancet* 2003; 361:296–300.
13. Cuzick J, Sestak I, Cawthorn S, et al. Tamoxifen for prevention of breast cancer: extended long-term follow-up of the IBIS-I breast cancer prevention trial. *Lancet Oncol* 2015; 16:67–75.
14. Fisher B, Costantino JP, Wickerham DL, et al. Tamoxifen for prevention of breast cancer: report of the National Surgical Adjuvant Breast and Bowel Project P-1 Study. *J Natl Cancer Inst* 1998; 90:1371–1388.
15. Chang J, Powles TJ, Ashley SE, et al. The effect of tamoxifen and hormone replacement therapy on serum cholesterol, bone mineral density and coagulation factors in healthy postmenopausal women participating in a randomised, controlled tamoxifen prevention study. *Ann Oncol* 1996; 7:671–675.
16. Herrington DM, Klein KP. Effects of SERMs on important indicators of cardiovascular health: lipoproteins, hemostatic factors and endothelial function. *Womens Health Issues* 2001; 11:95–102.
17. Kedar RP, Bourne TH, Powles TJ, et al. Effects of tamoxifen on uterus and ovaries of postmenopausal women in a randomized breast cancer prevention trial. *Lancet* 1994; 343:1318–1321.
18. Ettinger B, Black DM, Mitlak BH, et al. Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene: results from a 3-year randomized clinical trial. Multiple Outcomes of Raloxifene Evaluation (MORE) Investigators. *JAMA* 1999; 282:637–645.
19. Maricic M, Adachi JD, Sarkar S, Wu W, Wong M, Harper KD. Early effects of raloxifene on clinical vertebral fractures at 12 months in postmenopausal women with osteoporosis. *Arch Intern Med* 2002; 162:1140–1143.
20. Recker RR, Mitlak BH, Ni X, Krege JH. Long-term raloxifene for postmenopausal osteoporosis. *Curr Med Res Opin* 2011; 27:1755–1761.
21. Cummings SR, Eckert S, Krueger KA, et al. The effect of raloxifene on risk of breast cancer in postmenopausal women: results from the MORE randomized trial. Multiple Outcomes of Raloxifene Evaluation. *JAMA* 1999; 281:2189–2197.
22. Martino S, Cauley JA, Barrett-Connor E, et al; CORE Investigators. Continuing outcomes relevant to Evista: breast cancer incidence in postmenopausal osteoporotic women in a randomized trial of raloxifene. *J Natl Cancer Inst* 2004; 96:1751–1761.
23. Vogel VG, Qu Y, Wong M, Mitchell B, Mershon JL. Incidence of invasive breast cancer in

postmenopausal women after discontinuation of long-term raloxifene administration. *Clin Breast Cancer* 2009; 9:45–50.

24. Grady D, Cauley JA, Stock JL, et al. Effect of raloxifene on all-cause mortality. *Am J Med* 2010; 123:469.e1–e7.
25. Barrett-Connor E, Mosca L, Collins P, et al; Raloxifene Use for The Heart (RUTH) Trial Investigators. Effects of raloxifene on cardiovascular events and breast cancer in postmenopausal women. *N Engl J Med* 2006; 355:125–137.
26. Vogel VG. The NSABP Study of Tamoxifen and Raloxifene (STAR) trial. *Expert Rev Anticancer Ther* 2009; 9:51–60.
27. Vogel VG, Costantino JP, Wickerham DL, et al; National Surgical Adjuvant Breast and Bowel Project (NSABP). Effects of tamoxifen vs raloxifene on the risk of developing invasive breast cancer and other disease outcomes: the NSABP Study of Tamoxifen and Raloxifene (STAR) P-2 trial. *JAMA* 2006; 295:2727–2741.
28. Barnes KN, Pearce EF, Yancey AM, Forinash AB. Ospemifene in the treatment of vulvovaginal atrophy. *Ann Pharmacother* 2014; 48:752–757.
29. Rutanen EM, Heikkinen J, Halonen K, Komi J, Lammintausta R, Ylikorkala O. Effects of ospemifene, a novel SERM, on hormones, genital tract, climacteric symptoms, and quality of life in postmenopausal women: a double-blind, randomized trial. *Menopause* 2003; 10:433–439.
30. Constantine G, Graham S, Koltun WD, Kingsberg SA. Assessment of ospemifene or lubricants on clinical signs of VVA. *J Sex Med* 2014; 11:1033–1041.
31. Kingsberg SA, Wysocki S, Magnus L, Krychman ML. Vulvar and vaginal atrophy in postmenopausal women: findings from the REVIVE survey. *J Sex Med* 2013; 10:1790–1799.
32. Portman DJ, Bachmann GA, Simon JA; Ospemifene Study Group. Ospemifene, a novel selective estrogen receptor modulator for treating dyspareunia associated with postmenopausal vulvar and vaginal atrophy. *Menopause* 2013; 20:623–630.
33. Bachmann GA, Komi JO; Ospemifene Study Group. Ospemifene effectively treats vulvovaginal atrophy in postmenopausal women: results from a pivotal phase 3 study. *Menopause* 2010; 17:480–486.
34. Goldstein SR, Bachmann GA, Koninckx PR, Lin VH, Portman DJ, Ylikorkala O; Ospemifene Study Group. Ospemifene 12-month safety and efficacy in postmenopausal women with vulvar and vaginal atrophy. *Climacteric* 2014; 17:173–182.
35. Cui Y, Zong H, Yan H, Li N, Zhang Y. The efficacy and safety of ospemifene in treating dyspareunia associated with postmenopausal vulvar and vaginal atrophy: a systematic review

and meta-analysis. *J Sex Med* 2014; 11:487–497.

36. Komi J, Heikkinen J, Rutanen EM, Halonen K, Lammintausta R, Ylikorkala O. Effects of ospemifene, a novel SERM, on biochemical markers of bone turnover in healthy postmenopausal women. *Gynecol Endocrinol* 2004; 18:152–158.
37. Komi J, Lankinen KS, DeGregorio M, et al. Effects of ospemifene and raloxifene on biochemical markers of bone turnover in postmenopausal women. *J Bone Miner Metab* 2006; 24:314–318.
38. Wurz GT, Read KC, Marchisano-Karpman C, et al. Ospemifene inhibits the growth of dimethylbenzanthracene-induced mammary tumors in Sencar mice. *J Steroid Biochem Mol Biol* 2005; 97:230–240.
39. Pinkerton JV, Utian WH, Constantine GD, Olivier S, Pickar JH. Relief of vasomotor symptoms with the tissue-selective estrogen complex containing bazedoxifene/conjugated estrogens: a randomized, controlled trial. *Menopause* 2009; 16:1116–1124.
40. Pickar JH, Mirkin S. Tissue-selective agents: selective estrogen receptor modulators and the tissue-selective estrogen complex. *Menopause Int* 2010; 16:121–128.
41. Levine JP. Treating menopausal symptoms with a tissue-selective estrogen complex. *Gend Med* 2011; 8:57–68.
42. Lindsay R, Gallagher JC, Kagan R, Pickar JH, Constantine G. Efficacy of tissue-selective estradiol complex of bazedoxifene/conjugated estrogens for osteoporosis prevention in at-risk postmenopausal women. *Fertil Steril* 2009; 92:1045–1052.
43. Bachmann G, Bobula J, Mirkin S. Effects of bazedoxifene/conjugated estrogens on quality of life in postmenopausal women with symptoms of vulvar/vaginal atrophy. *Climacteric* 2010; 13:132–140.
44. Pickar JH, Yeh IT, Bachmann G, Speroff L. Endometrial effects of a tissue selective estrogen complex containing bazedoxifene/conjugated estrogens as a menopausal therapy. *Fertil Steril* 2009; 92:1018–1024.
45. Archer DF, Lewis V, Carr BR, Olivier S, Pickar JH. Bazedoxifene/conjugated estrogens (BZA/CE): incidence of uterine bleeding in postmenopausal women. *Fertil Steril* 2009; 92:1039–1044.
46. Pinkerton JV, Abraham L, Bushmakin AG, et al. Evaluation of the efficacy and safety of bazedoxifene/conjugated estrogens for secondary outcomes including vasomotor symptoms in postmenopausal women by years since menopause in the Selective estrogens, Menopause and Response to Therapy (SMART) trials. *J Womens Health (Larchmt)* 2014; 23:18–28.

Related Articles

REVIEWS

Is there a time limit for systemic menopausal hormone therapy?

The duration needs to be an individualized, shared decision that is assessed annually.



THIS PUBLICATION

[About CCJM](#), [Submit A Manuscript](#), [Author Information](#), [Subscription Information](#), [CME Calendar](#),
[Advertise With CCJM](#), [RSS Feeds](#), [Contact Us](#)

TOP SECTIONS

[CME](#), [Reviews](#), [1-Minute Consult](#), [Interpreting Key Trials](#), [IM Board Review](#), [Smart Testing](#)

HIGHLIGHTED FROM THIS SITE AND NETWORK

[About Cleveland Clinic](#), [Cleveland Clinic Education Institute](#),
[Cleveland Clinic Center for Continuing Education](#),
[Cleveland Clinic Resources for Medical Professionals](#),
[Cleveland Clinic Consumer Health Information](#), [Consult QD](#), [Contact ClinicalEdge Manager](#),
[Contact MD-IQ Manager](#), [MedJob Network](#)

[Advertise with MDedge](#)

See more with MDedge! [See our Other Publications](#)

MDedge: Your Source for Clinical Advantage

Copyright © 2017 Cleveland Clinic. All rights reserved. The information provided is for educational purposes only. Use of this website is subject to the [disclaimer](#) and [privacy policy](#).